# antibodies -online.com





anti-CLNK antibody (C-Term)



Overview

Clonality:

Image



Go to Product page

Quantity:	100 μL
Target:	CLNK
Binding Specificity:	C-Term
Reactivity:	Human, Rabbit, Cow, Dog, Guinea Pig, Horse, Mouse, Pig, Rat
Host:	Rabbit

Conjugate: This CLNK antibody is un-conjugated

Application: Western Blotting (WB)

Polyclonal

#### **Product Details**

Immunogen:	The immunogen is a synthetic peptide directed towards the C-terminal region of Human CLNK
Sequence:	SRQAVEEAFM KENKDGSFLV RDCSTKSKEE PYVLAVFYEN KVYNVKIRFL
Predicted Reactivity:	Cow: 79%, Dog: 86%, Guinea Pig: 79%, Horse: 86%, Human: 100%, Mouse: 93%, Pig: 86%, Rabbit: 100%, Rat: 86%
Characteristics:	This is a rabbit polyclonal antibody against CLNK. It was validated on Western Blot.
Purification:	Affinity Purified

## **Target Details**

Target:	CLNK
Alternative Name:	CLNK (CLNK Products)

## **Target Details**


Background:	MIST is a member of the SLP76 family of adaptors. MIST plays a role in the regulation of
	immunoreceptor signaling, including PLC-gamma-mediated B cell antigen receptor (BCR)
	signaling and FC-epsilon R1-mediated mast cell degranulation.
	Alias Symbols: MGC35197, MIST
	Protein Interaction Partner: ERBB2, EGFR, LCP2,
	Protein Size: 428
Molecular Weight:	49 kDa
Gene ID:	116449
NCBI Accession:	NM_052964, NP_443196
UniProt:	Q7Z7G1

# Application Details

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 428 AA
Restrictions:	For Research Use only

# Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.



#### **Western Blotting**

**Image 1.** WB Suggested Anti-CLNK Antibody Titration: 1.0 ug/ml Positive Control: Fetal Heart