

## Datasheet for ABIN2790984

## anti-CPSF3L antibody (C-Term)





Go to Product page

1	image	

Overview	
Quantity:	100 μL
Target:	CPSF3L
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Horse, Guinea Pig, Cow, Dog, Goat, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CPSF3L antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the C-terminal region of human CPSF3L
Sequence:	TESTYATTIR DSKRCRERDF LKKVHETVER GGKGAHEPEG AHLLLHGADR
Predicted Reactivity:	Cow: 100%, Dog: 100%, Goat: 77%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rat: 100%, Zebrafish: 100%
Characteristics:	This is a rabbit polyclonal antibody against CPSF3L. It was validated on Western Blot.
Purification:	Affinity Purified
Target Details	
Target:	CPSF3L

## Target Details

Restrictions:

Alternative Name:	CPSF3L (CPSF3L Products)	
Background:	The Integrator complex contains at least 12 subunits and associates with the C-terminal	
	domain of RNA polymerase II large subunit (POLR2A, MIM 180660) and mediates the 3-prime	
	end processing of small nuclear RNAs U1 (RNU1, MIM 180680) and U2 (RNU2, MIM 180690).	
	INTS11, or CPSF3L, is the catalytic subunit of the Integrator complex (Baillat et al., 2005	
	[PubMed 16239144]).[supplied by OMIM, Mar 2008]	
	Alias Symbols: RC68, INT11, RC-68, INTS11, CPSF73L	
	Protein Size: 278	
Molecular Weight:	30 kDa	
Gene ID:	54973	
NCBI Accession:	NM_001256456, NP_001243385	
UniProt:	Q5TA45	
Application Details		

Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Handling Advice:	Avoid repeat freeze-thaw cycles.	
Storage:	-20 °C	
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.	

For Research Use only

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

**Western Blotting** 

Image 1.