

Datasheet for ABIN2791909 anti-SFT2D2 antibody (N-Term)

1 Image

Alternative Name:



Go to Product page

Overview	
Quantity:	100 μL
Target:	SFT2D2
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Rabbit, Cow, Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SFT2D2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the N-terminal region of Human
	SFT2D2
Sequence:	LKKVLSGQDT EDRSGLSEVV EASSLSWSTR IKGFIACFAI GILCSLLGTV
Predicted Reactivity:	Cow: 86%, Guinea Pig: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against SFT2D2. It was validated on Western Blot.
Purification:	Affinity Purified
Target Details	
Target:	SFT2D2

SFT2D2 (SFT2D2 Products)

Target Details

Background:	SFT2D2 may be involved in fusion of retrograde transport vesicles derived from an endocytic
	compartment with the Golgi complex.
	Alias Symbols: UNQ512, dJ747L4.C1.2
	Protein Interaction Partner: UBC,
	Protein Size: 160
Molecular Weight:	1710
Moleculai Weight.	17 kDa
Gene ID:	375035

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeat freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

60 kDa__ 40 kDa__ 29 kDa__ 22 kDa__ 10 kDa__

Host: Rabbit

Target Name: SFT2D2

Sample Tissue: HepG2 Cell Lysate

Antibody Dilution: 1.0µg/ml

Western Blotting

Image1. Host:RabbitTargetName:SFT2D2SampleTissue:HumanHepG2WholeCellAntibodyDilution:1ug/ml