
antibodies .- online.com







anti-SDCCAG3 antibody (AA 218-435)



()	11/0	K\ /	iew
	\cup	ועוי	$I \cap VV$

Quantity:	100 μL
Target:	SDCCAG3
Binding Specificity:	AA 218-435
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SDCCAG3 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Purpose:	SDCCAG3 antibody [C1C3], 1 mg/ml
Immunogen:	Recombinant fragment corresponding to a region within amino acids 218-435 of SDCCAG3
Isotype:	IgG

Target Details

Target:	SDCCAG3
Alternative Name:	SDCCAG3 (SDCCAG3 Products)
Molecular Weight:	48 kDa
Gene ID:	10807

Application Details

Application Notes:	Tested Western blot
	Application Notes
	Suggested starting dilutions are as follows: WB: 1:500-1:3000. Not yet tested in other
	applications. Optimal working dilutions should be determined experimentally by the end
	user.Sample (50 µg of whole cell lysate) A: mouse brain 10 % SDS PAGE 107-10586 diluted at
	1:3000 SampleA: 30 µg of MOLT4 whole cell lysate 10 % SDS PAGE107-10586 diluted at 1:1000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Concentration: Buffer:	1 mg/mL 0.1M Tris, 0.1M Glycine, 10 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative.
Buffer:	0.1M Tris, 0.1M Glycine, 10 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative.
Buffer: Preservative:	0.1M Tris, 0.1M Glycine, 10 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative. Thimerosal (Merthiolate)
Buffer: Preservative:	0.1M Tris, 0.1M Glycine, 10 % Glycerol (pH 7). 0.01 % Thimerosal was added as a preservative. Thimerosal (Merthiolate) This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE