antibodies -online.com





anti-DYNC1LI1 antibody (AA 451-523) (Biotin)



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	DYNC1LI1
Binding Specificity:	AA 451-523
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DYNC1LI1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DNCLI1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	DYNC1LI1
Alternative Name:	DNCLI1 (DYNC1LI1 Products)

Target Details

Background:	Synonyms: LIC1, DNCLI1, Cytoplasmic dynein 1 light intermediate chain 1, Dynein light chain A,		
	DLC-A, Dynein light intermediate chain 1, cytosolic, DYNC1LI1		
	Background: Acts as one of several non-catalytic accessory components of the cytoplasmic		
	dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter		
	proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the		
	intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in		
	binding dynein to membranous organelles or chromosomes. Probably involved in the		
	microtubule-dependent transport of pericentrin. Is required for progress throuh the spindle		
	assembly checkpoint. The phosphorylated form appears to be involved in the selective removal		
	of MAD1L1 and MAD1L2 but not BUB1B from kinetochores.		
Gene ID:	51143		
UniProt:	Q9Y6G9		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		
Expiry Date:	12 months		