## antibodies -online.com





## anti-ARL6IP5 antibody (Alexa Fluor 594)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Alternative Name:

Background:

Quantity:	100 μL	
Target:	ARL6IP5 (Arl6ip5)	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARL6IP5 antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human JWA	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	ARL6IP5 (Arl6ip5)	

Synonyms: Addicsin, ADP ribosylation like factor 6 interacting protein 5, Aip 5, Aip 5, ARL 6

interacting protein 5, ARL6IP5 antibody Cytoskeleton related vitamin A responsive protein,

Dermal papilla derived protein 11, DERP 11, DERP11, Glutamate transporter EEAC 1 associated protein, Glutamate transporter EEAC1 associated protein, GTRAP3 18, Hp 22, Hp22, HSPC 127,

JWA (Arl6ip5 Products)

HSPC127, JM 5, JM5, JMX, PRA 1 domain family 3, PRA 2, PRA1 domain family 3, PRA2, PRAF 3, PRAF3, Prenylated Rab acceptor protein 2, Protein JWa, Putative MAPK activating protein PM27, PRAF3\_HUMAN.

Background: JWA is a four-transmembrane environmental responsive protein which binds to the CC chemokine recepor 5 (CCR5), a major co-receptor for human immunodeficiency virus (HIV). JWA is involved in environmental stress-responsive pathways in K562 cells, an erythroleukemia cell line derived from patients with chronic myeloid leukemia. Environmental stressors to K562 cells such as heat shock, a higher temperature than the ideal body temperature of the organism from which the cell line was derived, and oxidative stress, the production of oxygen-centered free radicals, regulate and increase the expres-sion of JWA. This response to environmental stressors suggests similiarity of JWA to heat shock protein 70 (HSP70), which is upregulated by heat stress and toxic chemicals.

Gene ID:

10550

Pathways:

Dicarboxylic Acid Transport

## **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	