# antibodies -online.com





# anti-LATS1 antibody (pThr1079) (Alexa Fluor 594)



Go to Product page

_			
$\cup$	)ve	rv	iew

Quantity:	100 μL
Target:	LATS1
Binding Specificity:	pThr1079
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LATS1 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human LATS1 around the phosphorylation site of Thr1079	
Isotype:	IgG	
Specificity:	This sequence will cross react with LATS2 if phosphorylated at Thr1041	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	LATS1	
Alternative Name:	LATS1 (LATS1 Products)	
Background:	Synonyms: wts, WARTS, Serine/threonine-protein kinase LATS1, Large tumor suppressor	
	homolog 1, WARTS protein kinase, h-warts, LATS1	
	Background: Negative regulator of YAP1 in the Hippo signaling pathway that plays a pivotal role	
	in organ size control and tumor suppression by restricting proliferation and promoting	
	apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and	
	STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates	
	LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and	
	inactivates YAP1 oncoprotein and WWTR1/TAZ. Phosphorylation of YAP1 by LATS1 inhibits its	
	translocation into the nucleus to regulate cellular genes important for cell proliferation, cell	
	death, and cell migration. Acts as a tumor suppressor which plays a critical role in maintenance	
	of ploidy through its actions in both mitotic progression and the G1 tetraploidy checkpoint.	
	Negatively regulates G2/M transition by down-regulating CDK1 kinase activity. Involved in the	
	control of p53 expression. Affects cytokinesis by regulating actin polymerization through	
	negative modulation of LIMK1. May also play a role in endocrine function.	
Gene ID:	9113	
UniProt:	095835	
Pathways:	Regulation of Actin Filament Polymerization, Maintenance of Protein Location	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months