
# antibodies -online.com





# anti-SH2B3 antibody (AA 161-260) (Alexa Fluor 594)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	SH2B3
Binding Specificity:	AA 161-260
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH2B3 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human LNK
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Monkey
Purification:	Purified by Protein A.

# **Target Details**

Target:	SH2B3
Alternative Name:	LNK (SH2B3 Products)
Background:	Synonyms: IDDM20,Lymphocyte adapter protein,Lymphocyte specic adapter

protein,Lymphocyte specic adapter protein Lnk,Lymphocyte-specic adapter protein Lnk,SH2B adapter protein 3,SH2B adaptor protein 3,SH2B3\_HUMAN,Signal transduction protein Lnk.

Background: APS (adapter molecule containing PH and SH2 domains), SH2-B and Lnk compose a family of adapter proteins, which contain a pleckstrin homology (PH) domain, an SH2 domain and a tyrosine phosphorylation site. Stimulation of B cell receptor (BCR) or T cell receptor (TCR) results in the phosphorylation of the immunoreceptor tyrosine-based activation motif (ITAM) of BCR, TCR and several substrates. APS, SH2-B and Lnk may bind to the ITAM domain of BCR and TCR. Lnk is tyrosine phosphorylated in response to TCR stimulation and APS has been shown to be tyrosine phosphorylated in response to BCR stimulation.

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions: For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months