
# antibodies -online.com





# anti-MAGEB2 antibody (AA 101-200) (AbBy Fluor® 594)



Go to Product page

$\sim$	
( )\/\	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	MAGEB2
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAGEB2 antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MAGEB2
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

# **Target Details**

Target:	MAGEB2
Alternative Name:	MAGEB2 (MAGEB2 Products)
Background:	Synonyms: Cancer/testis antigen 3.2, CT3.2, DAM6, DSS AHC critical interval MAGE superfamily

6, DSS/AHC critical interval gene, from MAGE superfamily, 6, MAGE B2 antigen, MAGE XP 2, Mage-b2, Mage-rs2, melanoma antigen family B, 2, Melanoma antigen, family B, 1, Melanoma associated antigen B2, Smage-2 protein, Smage2.

Background: This gene is a member of the MAGEB gene family. The members of this family have their entire coding sequences located in the last exon, and the encoded proteins show 50 to 68 % sequence identity to each other. The promoters and first exons of the MAGEB genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. This gene is localized in the DSS (dosage-sensitive sex reversal) critical region. It is expressed in testis and placenta, and in a significant fraction of tumors of various histological types. The MAGEB genes are clustered on chromosome Xp22-p21. [provided by RefSeq, Jul 2008].

Gene ID:

4113

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months