

[Go to Product page](#)

Datasheet for ABIN2804424

**anti-Myosin VI antibody (AA 1101-1294) (Alexa Fluor 594)**

## Overview

Quantity:	100 µL
Target:	Myosin VI (MYO6)
Binding Specificity:	AA 1101-1294
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Myosin VI antibody is conjugated to Alexa Fluor 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Myosin VI
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	Myosin VI (MYO6)
Alternative Name:	MYO6/Myosin VI ( <a href="#">MYO6 Products</a> )

## Target Details

Background:	<p>Synonyms: Deafness autosomal recessive 37, DFNA 22, DFNA22, DFNB 37, DFNB37, KIAA0389, MYO 6, Myo6, MYO6_HUMAN, Myosin 6, Myosin VI, Myosin-VI, Myosin6, MyosinVI, Unconventional myosin-6.</p> <p>Background: Myosin VI a molecular motor involved in intracellular vesicle and organelle transport, is the only Myosin motor that binds to the pointed end of Actin. This unique Myosin has only one light chain in the lever-arm domain and has highly irregular stepping with a wide range of step sizes, unlike that of other characterized Myosins. It associates with Clathrin-coated vesicles and disabled 2, indicating a role for Myosin VI in endocytosis. Mouse Myosin VI is expressed within the sensory hair cells of the cochlea. Human Myosin VI is mapped to the centromeric region of chromosome 6, a region that shows syntenic homology with the corresponding mouse chromosome 9 region, where the Snell's Waltzer mutation is located. The behavioral effects of the mouse Snell's Waltzer mutation are lack of responsiveness to sound, hyperactivity, head tossing and circling, due to the disorganization and fusing of stereocilia bundles within the inner ear. Defects of Myosin VI cause autosomal dominant nonsyndromic sensori-neural deafness in humans. Human Myosin VI is expressed in fetal cochlea and brain, as well as in adult brain.</p>
-------------	--

Pathways:	<a href="#">Sensory Perception of Sound</a> , <a href="#">Dicarboxylic Acid Transport</a> , <a href="#">Asymmetric Protein Localization</a>
-----------	---

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

---

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months