# antibodies .- online.com





# anti-NSG1 antibody (AA 1-100) (Alexa Fluor 594)



0 -	4.4	Prod	400		_
1-()	10	Pron	ויאוו	nan	$\boldsymbol{\omega}$
$\circ$	LO.	1 100	uot	Dud	

$\sim$	
( )\/\	rview
$\cup$	1 410 44

Quantity:	100 μL	
Target:	NSG1 (D4S234E)	
Binding Specificity:	AA 1-100	
Reactivity:	Mouse, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NSG1 antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NEEP21
Isotype:	IgG
Cross-Reactivity:	Monkey, Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	NSG1 (D4S234E)
Alternative Name:	NEEP21 (D4S234E Products)

#### **Target Details**

#### Background:

Synonyms: neuron-enriched endosomal 21 kDa protein, NSG1\_HUMAN.

Background: NEEP21, also known as Brain neuron cytoplasmic protein 1, NSG1 (neuron-specific proteins family member 1), P21 or D4S234E, is a single pass type II membrane protein belonging to the NSG family. It is highly expressed during neuronal maturation but its expression is downregulated in adult tissues. NEEP21 predominantly localizes to Rab 4-positive early endosomes in the somatodendritic neuronal compartment and is essential for proper receptor sorting and recycling in neurons. It associates with GRIP1 and GluR-2 and mediates the surface expression of GluR-2. When this interaction is interrupted, GluR-2 accumulates in early endosomes and leads to changes in evoked synaptic current properties. In addition, NEEP21 forms a complex with the SNARE protein, Syntaxin 13 (also known as Syntaxin 12), and participates in the recycling of transferrin receptors (TFRs) and NTR2 (neurotensin receptor

#### **Application Details**

IF(IHC-P) 1:50-200

2).

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months