



[Go to Product page](#)

Datasheet for ABIN2804637

anti-Nogo B Receptor antibody (AA 167-210) (Alexa Fluor 594)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | Nogo B Receptor (NUS1) |
| Binding Specificity: | AA 167-210 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Nogo B Receptor antibody is conjugated to Alexa Fluor 594 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human NGBR/Nogo B receptor |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | Nogo B Receptor (NUS1) |
| Alternative Name: | NGBR/Nogo B receptor (NUS1 Products) |
| Background: | Synonyms: C6orf68, NGBR, nuclear undecaprenyl pyrophosphate synthase 1 homolog, NUS1, |

Target Details

NGBR_HUMAN, Dehydrodolichyl diphosphate synthase complex subunit NUS1, Cis-prenyltransferase subunit NgBR, Nogo-B receptor.

Background: Nogo is an oligodendrocyte-specific member of the Reticulon family and is a component of CNS white matter that inhibits axon outgrowth, induces collapse of growth cones of chick dorsal root ganglion cells, and inhibits the spreading of 3T3 fibroblasts. Nogo is expressed by oligodendrocytes but not by Schwann cells and associates primarily with the endoplasmic reticulum. Nogo exists in three different splice forms, Nogo-A, -B and -C. NgBR (Nogo-B receptor), also known as nuclear undecaprenyl pyrophosphate synthase 1 homolog, is a 293 amino acid single-pass type I membrane protein that acts as a specific receptor for the amino-terminus of Nogo-B. Through this interaction, NgBR is involved in the regulation of vascular remodeling and angiogenesis. NgBR also enhances Niemann-Pick type C2 protein (NPC2) stabilization. Knockdown of NgBR mRNA leads to decreased NPC2 levels, which results in the hallmarks of NPC2 mutation: increased intracellular cholesterol accumulation and a loss of sterol sensing.

Gene ID: 116150

UniProt: [Q96E22](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months