antibodies -online.com





Datasheet for ABIN2804927

anti-DPH1 antibody (AA 251-350) (Alexa Fluor 594)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	DPH1
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DPH1 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human OVCA1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DPH1
Alternative Name:	OVCA1 (DPH1 Products)

Target Details

Background:	Synonyms: Candidate tumor suppressor in ovarian cancer 1, Diphthamide biosynthesis protein	
	1, Diphthamide biosynthesis protein 2 homolog like 1, Diphthamide biosynthesis protein 2 like,	
	Diptheria toxin resistance protein required for diphthamide biosynthesis Saccharomyces like 1,	
	Diptheria toxin resistance protein required for diphthamide biosynthesis like 1, DPH like 1 S.	
	cerevisiae, DPH like 1, DPH1, DPH1 homolog S. cerevisiae, DPH1 homolog, DPH2 like 1, DPH2L,	
	DPH2L1, HsDph1, Ovarian cancer associated gene 1, Ovarian cancer associated gene 1 protein,	
	Ovarian tumor suppressor candidate 1, DPH1_HUMAN.	
	Background: OVCA1 is required for the first step in the synthesis of diphthamide, a unique	
	posttranslationally modified histidine found only in translation elongation factor-2 (EEF2, MIM	
	130610). This modification is conserved from archaebacteria to humans and serves as the	
	target for ADP-ribosylation and inactivation of EEF2 by diphtheria toxin (DT) and Pseudomonas	
	exotoxin A. OVCA1 is one of several enzymes involved in synthesis of diphthamide in EEF2.	
Gene ID:	1801	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	