antibodies.com

Datasheet for ABIN2804963 anti-CLEC11A antibody (AbBy Fluor® 594)



Overview	
Quantity:	100 μL
Target:	CLEC11A
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLEC11A antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Pleckstrin/p47
lsotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	CLEC11A
Alternative Name:	p47 (CLEC11A Products)
Background:	Synonyms: p47, Platelet 47 kDa protein, Platelet and leukocyte C kinase substrate, Platelet P47, Platelet p47 protein, Pleckstrin, PLEK, PLEK_HUMAN, FLJ27168, OTTHUMP00000159962. Background: Platelet P47, also known as NSFL1C, UBX1, UBXD10 or UBXN2C, is a 370 amino acid protein that localizes to both the nucleus and the golgi apparatus (specifically to golgi

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2804963 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

stacks) and contains one SEP domain and one UBX domain. Functioning as part of a ternary
complex with VCP (a protein involved in the heterotypic fusion of transport vesicles with their
target membranes) and Syntaxin 5, p47 interacts with and reduces the ATPase activity of VCP
and is required for the fragmentation of golgi stacks during mitosis and for subsequent
reassembly of golgi stacks after mitosis. p47 is subject to phosphorylation during mitosis,
which inhibits p47-golgi interaction and is, therefore, required for proper golgi stack formation
and cisternal regrowth. Human p47 shares 89 % sequence identity with its mouse counterpart,
suggesting a conserved role between species. Multiple isoforms of p47 exist due to alternative
splicing events.
5341
IF(IHC-P) 1:50-200
For Research Use only
For Research Use only
For Research Use only Liquid
Liquid
Liquid 1 µg/µL
Liquid 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Liquid 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Liquid 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. ProClin
Liquid 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
Liquid 1 μg/μL Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.