

## Datasheet for ABIN2805132

## anti-PCDH17 antibody (AA 31-130) (AbBy Fluor® 594)



(	11	$\sim$	<b>~</b>	ı i	$\sim$	1 A I
U	١V	Н	rv	1	Н	V۷

Overview		
Quantity:	100 μL	
Target:	PCDH17	
Binding Specificity:	AA 31-130	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PCDH17 antibody is conjugated to AbBy Fluor® 594	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human PCDH17	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	
Target Details		
Target:	PCDH17	
Alternative Name:	PCDH17 (PCDH17 Products)	
Background:	Synonyms: PCDH68, C030033F14Rik, Gm78, PCD17_HUMAN, PCDH17, PCDH-17, PCDH 17,	

PCH68, Protocadherin 17, Protocadherin 68, Protocadherin-17, Protocadherin-68.

Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDH17 is a 1,159 amino acid single-pass type I membrane protein that contains six cadherin domains. Expressed as multiple alternatively spliced isoforms, PCDH17 is thought to function as a calcium-dependent cell adhesion protein that may play a role in establishing cell-cell connections within brain tissue. The gene encoding PCDH17 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4 % of the human genome.

## **Application Details**

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
Application Notes:	IF(IHC-P) 1:50-200

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months