

Datasheet for ABIN2805196

anti-PDGFA antibody (AA 102-213) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	PDGFA
Binding Specificity:	AA 102-213
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDGFA antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from rabbit PDGF-A
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PDGFA
Alternative Name:	PDGF A (PDGFA Products)

Target Details

Background:	Synonyms: PDGFA, PDGF AA, PDGF-AA, PDGF A chain, PDGF subunit A, PDGF-1, PDGF1, PDGFA, PDGFA_HUMAN, Platelet derived growth factor A chain precursor, Platelet derived growth factor alpha polypeptide, Platelet-derived growth factor A chain, Platelet-derived growth factor alpha polypeptide, Platelet-derived growth factor subunit A. Background: Platelet-derived growth factor is a potent mitogen for cells of mesenchymal origin. Binding of this growth factor to its affinity receptor elicits a variety of cellular responses. It is released by platelets upon wounding and plays an important role in stimulating adjacent cells to grow and thereby heal the wound. Antiparallel disulfide-linked dimer of nonidentical (A and B) chains. Homodimers of A and B chains are implicated in transformation processes. The long form contains a basic insert which acts as a cell retention signal. A-A and B-B, as well as A-B, dimers can bind to the PDGF receptor. It belongs to the PDGF/VEGF growth factor family.
Gene ID:	5154
UniProt:	P04085
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Smooth Muscle Cell Migration , Platelet-derived growth Factor Receptor Signaling

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months