

[Go to Product page](#)

Datasheet for ABIN2805245

anti-PEG10 antibody (AA 301-400) (Alexa Fluor 594)

Overview

Quantity:	100 µL
Target:	PEG10
Binding Specificity:	AA 301-400
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PEG10 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PEG10
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	PEG10
Alternative Name:	PEG10 (PEG10 Products)

Target Details

Background:	<p>Synonyms: Paternally expressed gene 10, MyEF 3, AA407948, Edr, Embryonal carcinoma differentiation regulated, HB 1, HB1, KIAA1051, Mar2, Mart2, MEF3 like 1, MEF3L, MEF3L1, Paternally expressed 10, Paternally expressed gene 10 ORF1, PEG10 protein, Putative uncharacterized protein PEG10, Retrotransposon gag domain containing 3, RGAG3, PEG10_HUMAN, Retrotransposon-derived protein PEG10, Embryonal carcinoma differentiation-regulated protein, Mammalian retrotransposon-derived protein 2, Myelin expression factor 3-like protein 1, MEF3-like protein 1, Paternally expressed gene 10 protein, Retrotransposon gag domain-containing protein 3, Retrotransposon-derived gag-like polyprotein, Ty3/Gypsy-like protein, EDR.</p> <p>Background: This is a paternally expressed imprinted gene that encodes transcripts containing two overlapping open reading frames (ORFs), RF1 and RF1/RF2, as well as retroviral-like slippage and pseudoknot elements, which can induce a -1 nucleotide frame-shift. ORF1 encodes a shorter isoform with a CCHC-type zinc finger motif containing a sequence characteristic of gag proteins of most retroviruses and some retrotransposons. The longer isoform is the result of -1 translational frame-shifting leading to translation of a gag/pol-like protein combining RF1 and RF2. It contains the active-site consensus sequence of the protease domain of pol proteins. Additional isoforms resulting from alternatively spliced transcript variants, as well as from use of upstream non-AUG (CUG) start codon, have been reported for this gene. Increased expression of this gene is associated with hepatocellular carcinomas. [provided by RefSeq, May 2010].</p>
-------------	---

Gene ID:	23089
----------	-------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months