
antibodies -online.com





Datasheet for ABIN2805314

anti-PHYHIP antibody (AA 231-330) (Alexa Fluor 594)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	PHYHIP
Binding Specificity:	AA 231-330
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PHYHIP antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PHYHIP
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PHYHIP
Alternative Name:	PHYHIP (PHYHIP Products)
Background: Synonyms: DYRK1A interacting protein 3, DYRK1AP3, PAHX AP1, PAHXAP1, phytanoyl	

hydroxylase interacting protein, phytanoyl CoA alpha hydroxylase associated protein, Phytanoyl CoA hydroxylase associated protein 1, Phytanoyl CoA hydroxylase interacting protein, Similar to a putative C.elegans gene encoded in cosmid M01B2, PHYIP_HUMAN.

Background: PHYHIP (Phytanoyl-CoA hydroxylase-interacting protein) is a 330 amino acid protein that is strongly expressed in brain, with weak expression in ovary, small intestine and ovary. In transgenic mice, overexpression of PHYHIP in heart results in tachycardia and tachyarrhythmia. PHYHIP interacts with the Refsum disease gene product, PAHX, indicating that PHYHIP may play a role in the CNS deficits of Refsum disease, which is characterized by cerebellar degeneration, neurologic damage and peripheral neuropathies. PHYHIP also interacts with Dyrk1A, a protein that that is overexpressed in brain of Down-syndrome patients, therefore PHYHIP may participate in some of the neurological abnormalities of Down syndrome. Significantly, the gene encoding PHYHIP is localized to a region of the short arm of human chromosome 8 that is frequently found deleted in prostate, breast and several other

Application Details

Application Notes:

types of cancers.

IF(IHC-P) 1:50-200

, .pp	(
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	