antibodies -online.com





anti-Prominin 2 antibody (AA 301-400) (Alexa Fluor 594)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	Prominin 2 (PROM2)
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prominin 2 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Prominin 2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig
Purification:	Purified by Protein A.

Target Details

Target:	Prominin 2 (PROM2)
Alternative Name:	Prominin 2 (PROM2 Products)

Target Details

Background:	Synonyms: PROML2, Prominin-2, PROM-2, Prominin-like protein 2, hPROML2, PROM2,	
	UNQ2521/PRO6014	
	Background: Prominin 2 is a 112 kDa glycoporotein structurally related to Prominin 1 (CD133)	
	although amino acid similarity is not more than 30 $\%$, but their genomic organization is	
	strikingly similar. Like Prominin 1, the prominin 2 exhibit similar membrane topology with 5	
	trans-membrane domains and two large glycosylated extracellular domains. Similar to	
	Prominin1 localization, the Prominin 2 is also associated with membrane protrusions of the	
	epithelial cells from adult kidney, and all along the digestive track and other epithelial	
	tissues.Prominin 2 expression is down-regulated in aggressive prostate cancer cell lines and	
	transient transfection of PROML2 expression vectors has been shown to induce apoptosis in	
	cultured prostate cancer cells, suggesting a tumor suppressive role for Prominin 2. Prominin 2	
	expression is likely to be involved in growth suppression in the prostate, and down-regulation of	
	Prominin 2 may disrupt normal prostatic homeostasis and lead to uncontrolled prostatic	
	growth.	
Gene ID:	150696	
UniProt:	Q8N271	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months