antibodies .- online.com





Datasheet for ABIN2805726

Overview

Target Details

Alternative Name:

Target:

RAB17

Rab17 (RAB17 Products)

anti-RAB17 antibody (AA 101-200) (AbBy Fluor® 594)

00	+~	Drad		page
()()	1()	Proa	uci	nade
\sim				P ~ 9 ~

Quantity:	100 μL	
Target:	RAB17	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RAB17 antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Rab17	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog	
Purification:	Purified by Protein A.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/2 | Product datasheet for ABIN2805726 | 06/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:

Synonyms: Rab 17, Rab17, RAB17 member RAS oncogene family, RAB17_HUMAN, Ras related protein Rab 17, Ras related protein Rab17, Ras-related protein Rab-17, FLJ12538, OTTHUMP00000164417, OTTHUMP00000202805.

Background: Rab17 belongs to the Rab family of small Ras-like GTPases. It is specifically expressed in epithelial cells and is upregulated during cell polarization. Immunofluorescence staining studies indicate that Rab17 is associated with the perinuclear recycling endosome in nonpolarized epithelial cells and with the apical recycling endosome in polarized epithelial cells. The function of Rab17 remains unclear. Reports of Rab17 colocalization with internalized IgA in the apical endosome suggest that it may regulate receptor-mediated transcytosis. Rab17 has also been shown to regulate melanocytic filopodia formation and melanosome trafficking. siRNA knockdown of Rab17 in melanoma cells induces melanosome accumulation in the cell periphery.

Application Details

Application	Notes:
-------------	--------

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months