# antibodies -online.com





# anti-Receptor Accessory Protein 1 antibody (AA 53-150) (Alexa Fluor 594)



#### Overview

Quantity:	100 μL	
Target:	Receptor Accessory Protein 1 (REEP1)	
Binding Specificity:	AA 53-150	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Receptor Accessory Protein 1 antibody is conjugated to Alexa Fluor 594	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

# Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human REEP1	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	Receptor Accessory Protein 1 (REEP1)
Alternative Name:	REEP1 (REEP1 Products)

#### **Target Details**

Background:

Synonyms: C2orf23, Chromosome 2 open reading frame 23, FLJ13110, Receptor accessory protein 1, Receptor expression-enhancing protein 1, Reep1, REEP1\_HUMAN, SPG31.

Background: Transport of G protein-coupled receptors (GPCRs) to the cell surface membrane is critical for receptor-ligand recognition. Mammalian GPCR odorant receptors (ORs), when heterologously expressed in cells, are poorly expressed on the cell surface. REEP1 (receptor expression-enhancing protein 1), is a 201 amino acid multi-pass mitochondrion membrane protein that belongs to the DP1 family. REEP1 interacts with odorant receptor proteins and may enhance the cell surface expression of odorant receptors. Mutations in the REEP1 gene are the third most common cause of hereditary spastic paraplegia (HSP) after spastin and atlastin gene mutations. Mutations in the REEP1 gene also cause spastic paraplegia autosomal dominant type 31, a neurodegenerative disorder. The REEP1 gene is conserved in chimpanzee, dog, cow, mouse, rat, chicken, zebrafish, A.thaliana and rice, and maps to human chromosome 2p11.2.

## **Application Details**

Anı	plication	Notes:
, \PI	piication	I VOLCO.

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months