antibodies -online.com





anti-RLIM antibody (AA 550-600) (AbBy Fluor® 594)



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Overview		
Quantity:	100 μL	
Target:	RLIM	
Binding Specificity:	AA 550-600	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RLIM antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human RNF12	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	RLIM	
Alternative Name:	RNF12 (RLIM Products)	
Background:	Synonyms: RNF12, NY-REN-43, E3 ubiquitin-protein ligase RLIM, LIM domain-interacting RING	

finger protein, RING finger LIM domain-binding protein, R-LIM, RING finger protein 12, Renal

carcinoma antigen NY-REN-43, RLIM

Background: E3 ubiquitin-protein ligase. Acts as a negative coregulator for LIM homeodomain transcription factors by mediating the ubiquitination and subsequent degradation of LIM cofactors LDB1 and LDB2 and by mediating the recruitment the SIN3a/histone deacetylase corepressor complex. Ubiquitination and degradation of LIM cofactors LDB1 and LDB2 allows DNA-bound LIM homeodomain transcription factors to interact with other protein partners such as RLIM. Plays a role in telomere length-mediated growth suppression by mediating the ubiquitination and degradation of TERF1. By targeting ZFP42 for degradation, acts as an activator of random inactivation of X chromosome in the embryo, a stochastic process in which one X chromosome is inactivated to minimize sex-related dosage differences of X-encoded genes in somatic cells of female placental mammals.

Gene ID:

51132

UniProt:

Q9NVW2

Application Details

Application Notes:

IF(IHC-P) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months