

Datasheet for ABIN2806282

anti-Separase antibody (pSer1073) (AbBy Fluor® 594)



Go to Product page

Overview

Quantity:	100 μL
Target:	Separase (ESPL1)
Binding Specificity:	pSer1073
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Separase antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human ESPL1 around the phosphorylation site of Ser1073
Isotype:	IgG
Cross-Reactivity:	Human
Cross-Reactivity: Purification:	Human Purified by Protein A.
•	
Purification:	

Target Details		
Background:	Synonyms: Caspase like protein ESPL1, ESP 1, ESP1, ESP-1, ESPL 1, ESPL1, ESPL-1, Extra	
	spindle poles like 1, Extra spindle poles like 1 protein, Separin, Similar to fission yeast cut1and	
	gene, SSE, Separase, ESPL1_HUMAN.	
	Background: Separase is a cysteine protease that is essential for mitotic progression by	
	separating sister chromatids. Each cell must receive one chromatid of every chromosome,	
	during mitosis. Cohesin plays an important role in cohering sister chromatids during the	
	prophase through anaphase stages of mitosis, making certain that genomic information is	
	replicated accurately. As the cellular division process continues, separase destroys cohesin by	
	means of cleavage, allowing the chromatids to separate and divide with the cell. Separase	
	activity is highly regulated. It not only cleaves cohesin at the onset of anaphase but also cleaves	
	itself, promoting downregulation of separase after anaphase. Should a human cell become an	
	aneuploid (one too many or too few chromatids), the embryo most likely will not survive. Should	
	the embryo survive, it will most likely develop severe birth defects or later develop malignant	
	cancers.	
Gene ID:	9700	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

ProClin

-20 °C

handled by trained staff only.

Preservative:

Storage:

Precaution of Use:

Storage Comment:

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months