

Datasheet for ABIN2806356

anti-INPP5D antibody (AbBy Fluor® 594)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL
Target:	INPP5D
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This INPP5D antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SHIP1	
Isotype:	IgG	
Cross-Reactivity:	Human, Monkey, Mouse	
Predicted Reactivity:	Rat,Cow,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	INPP5D	
Alternative Name:	SHIP1 (INPP5D Products)	
Background:	Synonyms: Inositol polyphosphate 5 phosphatase of 145 kDa, 4, 5-trisphosphate 5-phosphatase 1, hp51CN, hSHIP, Inositol polyphosphate 5 phosphatase 145 kDa, Inositol	

polyphosphate 5 phosphatase, Inositol polyphosphate 5 phosphatase D, Inositol polyphosphate-5-phosphatase of 145 kDa, INPP 5D, INPP5D, INPP5D protein, MGC104855, MGC142140, MGC142142, p150 ship, p150Ship, Phosphatidylinositol 3,4,5 trisphosphate 5 phosphatase 1, Phosphatidylinositol-3, SH2 containing inositol 5 phosphatase, SH2 containing inositol phosphatase isoform b, SH2 domain containing inositol 5' phosphatase 1, SH2 domain containing inositol phosphatase 1, SH2 domain-containing inositol phosphatase 1, SH2 domain-containing inositol phosphatase 1, SH2 domain-containing inositol-5"-phosphatase 1, SHIP 1, SHIP-1, SHIP1, SHIP1_HUMAN, Signaling inositol polyphosphate 5 phosphatase SIP 145, Signaling inositol polyphosphate 5 phosphatase SIP 145, SIP 145, SIP 145, SIP-145, SIP145.

Background: SHIP1 is a member of the inositol polyphosphate-5-phosphatase (INPP5) family and contains an N-terminal SH2 domain, an inositol phosphatase domain, and two C-terminal protein interaction domains. Expression of this protein is restricted to hematopoietic cells where its movement from the cytosol to the plasma membrane is mediated by tyrosine phosphorylation in response to multiple cytokine and B and T cell receptor activation. At the plasma membrane, the protein hydrolyzes the 5' phosphate from phosphatidylinositol (3,4,5)-trisphosphate and inositol-1,3,4,5-tetrakisphosphate, thereby affecting multiple signaling pathways. Overall the protein functions as a negative regulator of myeliod cell proliferation and survival.

Gene ID:

3635

Pathways:

TCR Signaling, BCR Signaling, Warburg Effect

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid
Concentration: $1 \,\mu\text{g}/\mu\text{L}$
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months