

[Go to Product page](#)

Datasheet for ABIN2806632
anti-SP011 antibody (Alexa Fluor 594)

Overview

| | |
|--------------|---|
| Quantity: | 100 µL |
| Target: | SP011 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SP011 antibody is conjugated to Alexa Fluor 594 |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human SP011 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | SP011 |
| Alternative Name: | SP011 (SP011 Products) |
| Background: | <p>Synonyms: Meiotic recombination protein SPO 11, Meiotic recombination protein SP011, MGC39953, SPO 11, SP011 meiotic protein covalently bound to DSB homolog, SP011 meiotic protein covalently bound to DSB like, SP011_HUMAN.</p> <p>Background: Spo11 is a type II topoisomerase that is thought to generate the chromosome</p> |

Target Details

breaks that initiate meiotic recombination. The Spo11 protein initiates meiotic recombination by generating DNA double-strand breaks (DSBs) and is required for meiotic synapsis in *S. cerevisiae*. The DSBs are located mostly in promoter regions, where the chromatin is in an open configuration, and cluster in domains along the chromosome. Expression of the Spo11 is detected mainly in the testis, in agreement with its predicted function in the initiation of meiotic recombination. Disruption of Spo11 leads to severe gonadal abnormalities from defective meiosis and results in infertility.

Gene ID: 23626

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months