antibodies -online.com





anti-SRD5A1 antibody (Alexa Fluor 594)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	ㄷ	٧	۷

Quantity:	100 μL
Target:	SRD5A1
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SRD5A1 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SRD5A1
Isotype:	IgG
Cross-Reactivity:	Cow, Human, Sheep
Predicted Reactivity:	Mouse,Rat,Dog,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	SRD5A1
Alternative Name:	SRD5A1 (SRD5A1 Products)
Background:	Synonyms: 3-oxo-5-alpha-steroid 4-dehydrogenase 1, SR type 1, Steroid 5-alpha-reductase 1, S5AR 1, S5A1_HUMAN.

Background: Steroid 5a-Reductase is an important enzyme in androgen physiology because it catalyzes the conversion of testosterone into the more potent 5a-dihydro-testosterone, which mediates androgen effects on target tissues. The enzyme exists as two isoforms: type 1, which is expressed mainly in the skin, and type 2, which is expressed mainly in the prostate. In cultured human skin cells, 5a-Reductase 1 shows heterogeneity of protein, and has different levels of transcriptional and translational expression. 5a-Reductase 1 is expressed in all portions of the hair follicle, whereas 5a-Reductase 2 is expressed only in mesenchymal portions. In addition, 5a-Reductase 1 is mainly expressed in human breast carcinoma and may play a role in the in situ production and actions of the potent androgen 5a-dihydrotestosterone, including inhibition of cancer cell proliferation in hormone-dependent human breast carcinoma. The 5a-Reductase-3a-hydroxysteroid dehydrogenase complex is present in the human brain, suggesting that the complex may be involved in the synthesis of neuroactive steroids or the catabolism of neurotoxic steroids.

Pathways:

Metabolism of Steroid Hormones and Vitamin D, Steroid Hormone Biosynthesis, Response to Growth Hormone Stimulus, C21-Steroid Hormone Metabolic Process

Application Details

Application Notes:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format: Concentration:	Liquid 1 μg/μL

IF(IHC-P) 1:50-200

 Concentration:
 1 μg/μL

 Buffer:
 Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

 Preservative:
 Sodium azide

 Precaution of Use:
 This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

 Storage:
 -20 °C

 Storage Comment:
 Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

 Expiry Date:
 12 months