

Datasheet for ABIN2807296

anti-Tuberin antibody (pSer939) (AbBy Fluor® 594)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Tuberin (TSC2)
Binding Specificity:	pSer939
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tuberin antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Tuberin around the phosphorylation site of Ser939
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Tuberin (TSC2)
Alternative Name:	Tuberin/TSC2 (TSC2 Products)

Target Details

Background:	Synonyms: Tuberinphospho S939, Tuberin phospho Ser939, p-Tuberin Ser939, FLJ43106, LAM, OTTHUMP00000158940, OTTHUMP00000198394, OTTHUMP00000198395, TSC2, TSC2_HUMAN, TSC4, Tuberin, Tuberous sclerosis 2, Tuberous sclerosis 2 protein, Tuberous sclerosis 2 protein homolog. Background: Tuberin, or TSC2 (Tuberous sclerosis complex), is implicated as a tumor suppressor. It may function in vesicular transport, and may also play a role in the regulation of cell growth arrest and in the regulation of transcription mediated by steroid receptors. Interaction between hamartin (TSC1) and tuberin may facilitate vesicular docking. It specifically stimulates the intrinsic GTPase activity of the Ras related protein RAP1A and RAB5, suggesting a possible mechanism for its role in regulating cellular growth. Mutations in tuberin lead to constitutive activation of RAP1A in tumors. At least three isoforms of Tuberin exist.
Gene ID:	7249
Pathways:	RTK Signaling , AMPK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway , Regulation of Cell Size , Tube Formation , Protein targeting to Nucleus

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months