



[Go to Product page](#)

Datasheet for ABIN2807528

anti-WASF1 antibody (AA 481-559) (AbBy Fluor® 594)

Overview

Quantity:	100 µL
Target:	WASF1
Binding Specificity:	AA 481-559
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WASF1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human WAVE 1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	WASF1
Alternative Name:	WAVE 1 (WASF1 Products)

Target Details

Background: Synonyms: Protein WAVE-1, Protein WAVE1, SCAR1, Similar to a plant extensin like protein, Verprolin homology domain containing protein 1, Verprolin homology domain-containing protein 1, WAS protein family, member 1, Wasf1, WASF1_HUMAN, WASL, WASP family 1, WASP family protein member 1, WAVE, WAVE1, Wiskott Aldrich syndrome protein family member 1, Wiskott-Aldrich syndrome protein family member 1.

Background: WASP (for Wiskott-Aldrich syndrome protein) and N-WASP are downstream effectors of Cdc42 that are implicated in Actin polymerization and cytoskeletal organization. The WASP family also includes VASP (vasodilator-stimulated phosphoprotein) and Mena (for mammalian enabled protein), which accumulate at focal adhesions and are also involved in the regulation of the Actin cytoskeleton. The WAVE proteins are related to the WASP family proteins and are likewise involved in mediating Actin reorganization downstream of the Rho family of small GTPases. The protein homologs WAVE1 and WAVE2 regulate membrane ruffling by inducing the formation of Actin filament clusters in response to GTP binding and by activating Rac. They mediate Actin polymerization by cooperating with the Arp2/3 complex, thereby promoting the formation of Actin filaments. WAVE1, which is also designated SCAR (suppressor of cAR), is expressed primarily in the brain, while WAVE2 is widely expressed, with the expression highest in peripheral blood leukocytes. WAVE3 forms a multiprotein complex that links receptor kinases with Actin and plays a role in the transduction of signals involving changes in cell shape, function or motility.

Gene ID: 8936

Pathways: [RTK Signaling, Regulation of Actin Filament Polymerization](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months