# antibodies .- online.com





Datasheet for ABIN2808549

# anti-GM1123 antibody (AA 21-120) (AbBy Fluor® 594)



Go to Product page

( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	GM1123	
Binding Specificity:	AA 21-120	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GM1123 antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Coxsackie Adenovirus Receptor
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	GM1123
Alternative Name:	Coxsackie Adenovirus Receptor (GM1123 Products)

## **Target Details**

Storage Comment:

rarger betane		
Background:	Synonyms: CAR, HCAR, CAR4/6, Coxsackievirus and adenovirus receptor, CVB3-binding protein,	
	Coxsackievirus B-adenovirus receptor, HCVADR, CXADR	
	Background: Component of the epithelial apical junction complex that may function as an	
	homophilic cell adhesion molecule and is essential for tight junction integrity. Also involved in	
	transepithelial migration of leukocytes through adhesive interactions with AMICA1/JAML a	
	transmembrane protein of the plasma membrane of leukocytes. The interaction between both	
	receptors also mediates the activation of gamma-delta T-cells, a subpopulation of T-cells	
	residing in epithelia and involved in tissue homeostasis and repair. Upon epithelial CXADR-	
	binding, AMICA1 induces downstream cell signaling events in gamma-delta T-cells through PI3-	
	kinase and MAP kinases. It results in proliferation and production of cytokines and growth	
	factors by T-cells that in turn stimulate epithelial tissues repair.	
Gene ID:	1525	
UniProt:	P78310	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

1	1 11	١٠
$\vdash$	land	∥n∩
	ıarıa	

Expiry Date:

12 months