



[Go to Product page](#)

Datasheet for ABIN2808627

## anti-GABRA6 antibody (AA 85-180) (Alexa Fluor 594)

### Overview

Quantity:	100 µL
Target:	GABRA6
Binding Specificity:	AA 85-180
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABRA6 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GABRA6/GABA A Receptor alpha 6
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	GABRA6
Alternative Name:	GABRA6/GABA A Receptor alpha 6 ( <a href="#">GABRA6 Products</a> )
Background:	Synonyms: GABA A, GABA A Receptor alpha 6 polypeptide, GABA A receptor alpha 6, GABA A

## Target Details

---

receptor subunit alpha 6, GABA subunit A receptor alpha 6, GABAA receptor subunit alpha-6, GABRA 6, GABRA6, Gamma aminobutyric acid A receptor alpha 6, Gamma aminobutyric acid GABA A receptor alpha 6, Gamma aminobutyric acid receptor subunit alpha 6, Gamma-aminobutyric acid receptor subunit alpha-6, GBRA6\_HUMAN, MGC116903, MGC116904.

Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (gamma-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl<sup>-</sup> conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA R gamma 1-3, GABAA R gamma , GABAA R delta. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

---

Gene ID: 2559

---

UniProt: [Q16445](#)

---

Pathways: [Synaptic Membrane](#)

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

---

## Handling

---

handled by trained staff only.

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months