# antibodies -online.com





## anti-GABRG3 antibody (Alexa Fluor 594)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:    | 100 μL  |
|--------------|---|
| Target:      | GABRG3  |
| Reactivity:  | Human, Mouse, Rat                                     |
| Host:        | Rabbit  |
| Clonality:   | Polyclonal  |
| Conjugate:   | This GABRG3 antibody is conjugated to Alexa Fluor 594 |
| Application: | Western Blotting (WB)                                 |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human GABRG3/GABA A Receptor gamma 3 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat  |
| Purification:         | Purified by Protein A.   |

### Target Details

| Target:           | GABRG3  |  |
|-------------------|---|--|
| Alternative Name: | GABRG3/ GABA A Receptor gamma 3 (GABRG3 Products)                                   |  |
| Background:       | Synonyms: GABAA receptor, GABAA receptor subunit gamma 3, GABR G3, GABRG 3, GABRG3  |  |
|                   | Gamma aminobutyric acid GABA A receptor gamma 3, Gamma aminobutyric acid A receptor |  |

gamma 3, Gamma aminobutyric acid receptor subunit gamma 3, GBRG3\_HUMAN.

Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (gamma-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a CI- conductance that inhibits neuronal firing.

GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R Alpha 1-6, GABAA R Beta 1-3, GABAA R Gamma 1-3, GABAA R Delta, GABAA R Epsilon, GABAA R Zeta 1 and GABAA R Zeta 2. The GABAB family is composed of GABAB R1 Alpha and GABAB R1 Beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

#### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200   |
|--------------------|--|
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.             |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                      |
| Expiry Date:       | 12 months  |