

Datasheet for ABIN2809067

anti-TAP1 antibody (AA 501-600) (AbBy Fluor® 594)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TAP1
Binding Specificity:	AA 501-600
Reactivity:	Human, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAP1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Tap1/ABCB2
Isotype:	IgG
Cross-Reactivity:	Human, Pig
Predicted Reactivity:	Mouse,Rat,Cow,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	TAP1
Alternative Name:	Tap1 (TAP1 Products)

Target Details

Background:	Synonyms: APT1, PSF1, ABC17, ABCB2, PSF-1, RING4, TAP1N, D6S114E, TAP1*12N, Antigen peptide transporter 1, ATP-binding cassette sub-family B member 2, Peptide supply factor 1, Peptide transporter PSF1, Peptide transporter TAP1, Peptide transporter involved in antigen processing 1, Really interesting new gene 4 protein, TAP1, Y3 Background: Involved in the transport of antigens from the cytoplasm to the endoplasmic reticulum for association with MHC class I molecules. Also acts as a molecular scaffold for the final stage of MHC class I folding, namely the binding of peptide. Nascent MHC class I molecules associate with TAP via tapasin. Inhibited by the covalent attachment of herpes simplex virus ICP47 protein, which blocks the peptide-binding site of TAP. Inhibited by human cytomegalovirus US6 glycoprotein, which binds to the luminal side of the TAP complex and inhibits peptide translocation by specifically blocking ATP-binding to TAP1 and prevents the conformational rearrangement of TAP induced by peptide binding. Inhibited by human adenovirus E3-19K glycoprotein, which binds the TAP complex and acts as a tapasin inhibitor, preventing MHC class I/TAP association. Expression of TAP1 is down-regulated by human Epstein-Barr virus vIL-1 protein, thereby affecting the transport of peptides into the endoplasmic reticulum and subsequent peptide loading by MHC class I molecules.
Gene ID:	6890
UniProt:	Q03518
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Human Leukocyte Antigen (HLA) in Adaptive Immune Response

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months