



[Go to Product page](#)

Datasheet for ABIN2809300

## anti-5HT1D antibody (AA 2-110) (AbBy Fluor® 594)

### Overview

Quantity:	100 µL
Target:	5HT1D (HTR1D)
Binding Specificity:	AA 2-110
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This 5HT1D antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human 5HT1D Receptor/SR-1D
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	5HT1D (HTR1D)
Alternative Name:	5HT1D Receptor/SR-1D ( <a href="#">HTR1D Products</a> )

## Target Details

---

**Background:** Synonyms: 5-hydroxytryptamine receptor 1D, 5-HT-1D, 5-HT1D, Serotonin 1D alpha receptor, 5-HT-1D-alpha, Serotonin receptor 1D

Background: The members of the G-protein-coupled receptor family are distinguished by their slow transmitting response to ligand binding. These seven transmembrane proteins include the adrenergic, serotonin and dopamine receptors. The effect of the signaling molecule can be excitatory or inhibitory depending on the type of receptor to which it binds. b-adrenergic bound to adrenaline activates adenylyl cyclase, while a2-adrenergic receptor bound to adrenaline inhibits adenylyl cyclase. Like the a2-adrenergic receptor, serotonin receptor functions are also mediated by G proteins that inhibit the activity of adenylyl cyclase. The serotonin receptors have been classified into several categories, designated SR-17 (5HT17). Subtypes within the SR-1 group include SR-1A, -1B, -1D, -1E and -1F.

---

**Gene ID:** 3352

---

**UniProt:** [P28221](#)

---

**Pathways:** [JAK-STAT Signaling](#)

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 µg/µL

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

**Precaution of Use:** This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

**Storage:** -20 °C

---

**Storage Comment:** Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

## Handling

---

Expiry Date: 12 months