antibodies -online.com





Datasheet for ABIN2809799

anti-APIP antibody (AA 151-242) (Alexa Fluor 594)



| () | ve | K\ / | | A . |
|-----|--------|------|----|-----|
| | \cup | 1 V/ | Щ. | V۷ |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | APIP |
| Binding Specificity: | AA 151-242 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This APIP antibody is conjugated to Alexa Fluor 594 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human APIP |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | APIP |
|-------------------|---|
| Alternative Name: | Apaf1 Interacting Protein (APIP Products) |
| Background: | Synonyms: MTNB_HUMAN, Apaf1 Interacting Protein, APIP2, CG129, CGI 29, MMRP19, MTRu 1 |

| P dehydratase, Probable methylthioribulose 1 phosphate dehydrata |
|--|
|--|

Background: The mammalian homologues of the key cell death gene CED 4 in C. elegans has been identified recently from human and mouse and designated Apaf1 (for apoptosis protease activating factor 1). Apaf1 binds to cytochrome c (Apaf2) and caspase 9 (Apaf3), which leads to caspase 9 activation. Activated caspase 9 in turn cleaves and activates caspase 3 that is one of the key proteases, being responsible for the proteolytic cleavage of many key proteins in apoptosis. A new Apaf1 Interacting Protein (APIP) also known as CG129 and MMRP19, has been identified as a negative regulator of ischemic injury. APIP competes with Caspase 9 binding site of Apaf1. APIP is predicted to code for a 204 amino acid. An isoform of APIP, APIP2 encodes a 242 amino acid protein, which is an alternative splicing variant differing in its N terminus from APIP. APIP transcript is ubiquitously expressed in most adult tissue with high expression in skeletal muscle, heart, and kidney.

Pathways: Methionine Biosynthetic Process

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |

| | | 1. | |
|----------|-------|------|---|
| \vdash | land | lın | |
| | ıaııu | 1111 | U |
| | | | |

Expiry Date:

12 months