

Datasheet for ABIN2810274

anti-Abeta 1-42 antibody (AA 1-42) (AbBy Fluor® 594)



()	ve	r\/i	۱۸/
\cup	V C	1 / 1	 v v

Overview		
Quantity:	100 μL	
Target:	Abeta 1-42	
Binding Specificity:	AA 1-42	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Abeta 1-42 antibody is conjugated to AbBy Fluor® 594	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Flow Cytometry (FACS)	
Product Details		
Immunogen:	KLH conjugated synthetic peptide of human beta-Amyloid(1-42)	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Prodicted Popotivity:	Dog Cow Dig Chickon Pabhit	

Predicted Reactivity: Dog,Cow,Pig,Chicken,Rabbit

Purification: Purified by Protein A.

Target Details

Abeta 1-42 Target: Alternative Name: beta Amyloid 1-42 (Abeta 1-42 Products)

Target Details

Background:	Synonyms: AAA, AD1, PN2, ABPP, APPI, CVAP, ABETA, PN-II, CTFgamma, Amyloid beta A4		
	protein, APP, Alzheimer disease amyloid protein, Cerebral vascular amyloid peptide, PreA4,		
	Protease nexin-II, A4		
	Background: The cerebral and vascular plaques associated with Alzheimer's disease are mainly		
	composed of Amyloid beta peptides. beta Amyloid is derived from cleavage of the Amyloid		
	precursor protein and varies in length from 39 to 43 amino acids. beta Amyloid [1-40], beta		
	Amyloid [1-42], and beta Amyloid [1-43] peptides result from cleavage of Amyloid precursor		
	protein after residues 40, 42, and 43, respectively. The cleavage takes place by gamma-		
	secretase during the last Amyloid precursor protein processing step. beta Amyloid [1-40], beta		
	Amyloid [1-42], and beta Amyloid [1-43] peptides are major constituents of the plaques and		
	tangles that occur in Alzheimer's disease. beta Amyloid and peptides have been developed as		
	tools for elucidating the biology of Alzheimer's disease.		
Gene ID:	351		
UniProt:	P05067		
Application Details			
Application Notes:	FCM 1:20-100		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

	_	
-	II	1:
-	-222	nnc
-	Hand	1111()

Expiry Date:

12 months