antibodies -online.com





anti-BRP44 antibody (AA 21-100) (AbBy Fluor® 594)



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	BRP44
Binding Specificity:	AA 21-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRP44 antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BRP44
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	BRP44
Alternative Name:	BRP44 (BRP44 Products)
Background:	Synonyms: ESTM43, MGC125753, BR44_HUMAN, Brain protein 44, BRP44, DKFZp564B167,

MGC125752, rCG_46381, RGD1563422, RP1-295C6.1.

Background: BRP44 is a 127 amino acid protein belonging to the UPF0041 family. The gene that encodes BRP44 maps to human chromosome 1, which is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.

Gene ID:	25874

Pathways: Warburg Effect

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months