# antibodies -online.com





Go to Product page

### Datasheet for ABIN2810702

## anti-AIDA antibody (AbBy Fluor® 594)

|                 | Go to Product page  |
|-----------------|---|
| Overview        |   |
| Quantity:       | 100 μL  |
| Target:         | AIDA  |
| Reactivity:     | Human, Mouse, Rat   |
| Host:           | Rabbit  |
| Clonality:      | Polyclonal  |
| Conjugate:      | This AIDA antibody is conjugated to AbBy Fluor® 594                             |
| Application:    | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details |   |
| Immunogen:      | KLH conjugated synthetic peptide derived from human C1orf180/AIDA               |
| Isotype:        | IgG   |

| Immunogen:        | KLH conjugated synthetic peptide derived from human C1orf180/AIDA |
|-------------------|---|
| Isotype:          | IgG   |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Purified by Protein A.  |

#### **Target Details**

| Target:           | AIDA   |
|-------------------|--|
| Alternative Name: | C1orf80 (AIDA Products)  |
| Background:       | Synonyms: dorsalization-associated protein, AIDA, AIDA_HUMAN, Axin interaction partner and       |
|                   | dorsalization antagonist, Axin interactor, Axin interactor, dorsalization associated, Chromosome |
|                   | 1 open reading frame 80, UPF0491 protein C1orf80.  |
|                   | Background: AIDA, also known as C1orf80, is a 306 amino acid protein that belongs to the axin    |

#### Target Details

interactor family. Expressed in a variety of tissues, including skeletal muscle and heart, AIDA functions as a ventralizing factor during embryogenesis, disrupting Axin homodimerization and inhibiting Axin-mediated JNK activation. Axin, a scaffold protein, is important for both JNK signaling and the canonical Wnt pathway, two processes that play an essential role in embryonic dorsoventral patterning. Disruption of Axin by AIDA results in the negative regulation of JNK and Wnt signaling, thereby affecting embryonic developmental events. Three isoforms of AIDA exist due to alternative splicing events.

Gene ID:

439927

#### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200    |
|--------------------|-----------------------|
| Restrictions:      | For Research Use only |
| Handling           |                       |
| Format:            | Liquid                |
| Concentration:     | 1 μg/μL               |

| Concentration:     | 1 μg/μL  |
|--------------------|--|
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |