



[Go to Product page](#)

Datasheet for ABIN2811202

## anti-CCDC138 antibody (AA 551-665) (Alexa Fluor 594)

### Overview

Quantity:	100 µL
Target:	CCDC138
Binding Specificity:	AA 551-665
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC138 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC138
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	CCDC138
Alternative Name:	CCDC138 ( <a href="#">CCDC138 Products</a> )

## Target Details

Background:	<p>Synonyms: CCDC 138, CCDC138, Coiled coil domain containing 138, Coiled coil domain containing protein 138, coiled-coil domain containing 138, FLJ 32745, FLJ32745, CC138_HUMAN.</p> <p>Background: CCDC138, also known as FLJ32745, is a 685 amino acid protein expressed as two isoforms produced by alternative splicing. The gene that encodes CCDC138 maps to human chromosome 2q12.3. The second largest human chromosome, chromosome 2 consists of 237 million bases, encodes over 1,400 genes and makes up approximately 8 % of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alstr syndrome is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes.</p>
-------------	--

Gene ID:	165055
----------	--------

Pathways:	<a href="#">BCR Signaling</a>
-----------	-------------------------------

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months