antibodies -online.com





anti-CDC14B antibody (AA 65-160) (Alexa Fluor 594)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | CDC14B | |
| Binding Specificity: | AA 65-160 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CDC14B antibody is conjugated to Alexa Fluor 594 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CDC14B | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse, Rat | |
| Predicted Reactivity: | Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | CDC14B | |
|-------------------|--------------------------|--|
| Alternative Name: | CDC14B (CDC14B Products) | |

Target Details

| _ | | | | |
|---------------------|--|--|--|--|
| Background: | Synonyms: CDC14B3, Cdc14B1, Cdc14B2, hCDC14B, Dual specificity protein phosphatase | | | |
| | CDC14B, CDC14 cell division cycle 14 homolog B, CDC14B | | | |
| | Background: Dual-specificity phosphatase involved in DNA damage response. Essential | | | |
| | regulator of the G2 DNA damage checkpoint: following DNA damage, translocates to the | | | |
| | nucleus and dephosphorylates FZR1/CDH1, a key activator of the anaphase promoting complex/cyclosome (APC/C). Dephosphorylates SIRT2 around early anaphase. | | | |
| | | | | |
| | Dephosphorylation of FZR1/CDH1 activates the APC/C, leading to the ubiquitination of PLK1, | | | |
| | preventing entry into mitosis. Preferentially dephosphorylates proteins modified by proline- | | | |
| | directed kinases. | | | |
| Gene ID: | 8555 | | | |
| UniProt: | 060729 | | | |
| Application Details | | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | | |
| | IF(IHC-F) 1:50-200 | | | |
| | IF(ICC) 1:50-200 | | | |
| Restrictions: | For Research Use only | | | |
| Handling | | | | |
| Format: | Liquid | | | |
| Concentration: | 1 μg/μL | | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | | |
| | 50 % Glycerol. | | | |
| Preservative: | ProClin | | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | | |
| | handled by trained staff only. | | | |
| Storage: | -20 °C | | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | | |
| Expiry Date: | 12 months | | | |
| | | | | |