



[Go to Product page](#)

Datasheet for ABIN2811462

anti-FAT4 antibody (AA 4351-4450) (AbBy Fluor® 594)

Overview

Quantity:	100 µL
Target:	FAT4
Binding Specificity:	AA 4351-4450
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAT4 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDHF14
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	FAT4
Alternative Name:	CDHF14 (FAT4 Products)

Target Details

Background: Synonyms: Cadherin family member 14, CDHF14, FAT tumor suppressor homolog 4, Fat-like cadherin protein FAT-J, FAT4, FAT4_HUMAN, FATJ, hFat4, Nbla00548, Protocadherin Fat 4.

Background: The cadherins represent a family of Ca²⁺-dependent adhesion molecules that function to mediate cell to cell binding that is critical for the maintenance of structure and morphogenesis. Cadherins contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short C-terminal intracellular domain interacts with a variety of cytoplasmic proteins, including beta-catenin, to regulate cadherin function. The cadherin superfamily includes cadherins, protocadherins, desmogleins and desmocollins. FAT4 (FAT tumor suppressor homolog 4), also known as FAT-J, CDHF14 or CDHR11, is a 4,981 amino acid single-pass type I membrane protein that belongs to the protocadherin subfamily of cadherins and localizes to the primary cilia of kidney. Widely expressed, FAT4 contains thirty-four cadherin domains, six EGF-like domains and two laminin G-like domains. FAT4 may function in the regulation of planar cell polarity.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months