

Datasheet for ABIN2811478

anti-CDK5RAP3 antibody (AA 51-150) (AbBy Fluor® 594)



Go to Product page

_				
()	VE	r\/		Λ/
()	V C	I V	1	v v

Overview	
Quantity:	100 μL
Target:	CDK5RAP3
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK5RAP3 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human CDK5RAP3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep
Purification:	Purified by Protein A.

Target Details

Target:	CDK5RAP3
Alternative Name:	CDK5RAP3 (CDK5RAP3 Products)
Background:	Synonyms: C53, CDK5 activator binding protein C53, CDK5 regulatory subunit associated

Storage Comment:

12 months

Expiry Date:

rarget Details	
	protein 3, HSF 27, HSF27, IC53, Ischemic heart CDK5 activator binding protein C53,
	LXXLL/leucine zipper containing ARFbinding protein, LZAP, MST016, OK/SW cl.114, PP1553,
	CK5P3_HUMAN.
	Background: Neuronal CDC2-like kinase, which is involved in the regulation of neuronal
	differentiation, is composed of a catalytic subunit, CDK5, and an activating subunit, p25NCK5A.
	The protein encoded by this gene binds to p25NCK5A and therefore may be involved in
	neuronal differentiation. The encoded protein, which may be a substrate of neuronal CDC2-like
	kinase, has also been found in vascular endothelial cells, where it mediates cell proliferation.
Gene ID:	80279
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.