

[Go to Product page](#)

Datasheet for ABIN2811560

anti-AGAP2 antibody (AA 981-1080) (Alexa Fluor 594)

Overview

Quantity:	100 µL
Target:	AGAP2
Binding Specificity:	AA 981-1080
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AGAP2 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Centaurin gamma 1
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	AGAP2
Alternative Name:	Centaurin gamma 1 (AGAP2 Products)

Target Details

Background:	<p>Synonyms: Centaurin gamma 1, ANK repeat and PH domain-containing protein 2, AGAP-2, Agap2, Agap2 ArfGAP with GTPase domain, ankyrin repeat and PH domain 2, AGAP2_HUMAN, ANK repeat and PH domains 2, Arf GAP with GTP binding protein like, Arf GAP with GTP-binding protein-like, ANK repeat and PH domains 2, Arf GAP with GTP-binding protein-like, ankyrin repeat and pleckstrin homology domains 2, Arf-GAP with GTPase, Arf-GAP with GTPase, ANK repeat and PH domain-containing protein 2, ArfGAP with GTPase domain, ankyrin repeat and PH domain 2, centaurin gamma1, Centaurin-gamma-1, Centg1, cnt g1, Cnt-g1, Cnt2, GGAP2, GTP binding and GTPase activating protein 2, GTP-binding and GTPase activating protein 2, GTP-binding and GTPase-activating protein 2, Nuclear GTPase PIKE, Phosphatidylinositol-3-kinase enhancer, Phosphoinositide 3 kinase enhancer, PI3K ENHANCER, PIKE.</p> <p>Background: GTPase-activating protein (GAP) for ARF1 and ARF5, which also shows strong GTPase activity. Isoform 1 participates in the prevention of neuronal apoptosis by enhancing PI3 kinase activity. It aids the coupling of metabotropic glutamate receptor 1 (GRM1) to cytoplasmic PI3 kinase by interacting with Homer scaffolding proteins, and also seems to mediate anti-apoptotic effects of NGF by activating nuclear PI3 kinase. Isoform 2 does not stimulate PI3 kinase but may protect cells from apoptosis by stimulating Akt. It also regulates the adapter protein 1 (AP-1)-dependent trafficking of proteins in the endosomal system. It seems to be oncogenic. It is overexpressed in cancer cells, prevents apoptosis and promotes cancer cell invasion.</p>
-------------	--

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months