antibodies -online.com





anti-CREB3L3 antibody (AA 201-300) (Alexa Fluor 594)



| Go to | D | | |
|-------|------|-----|------|
| | Pron | HCL | nane |
| | | | |

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | CREB3L3 |
| Binding Specificity: | AA 201-300 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CREB3L3 antibody is conjugated to Alexa Fluor 594 |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CREB-H |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CREB3L3 |
|-------------------|-----------------------------------|
| Alternative Name: | CREB-H/CREB3L3 (CREB3L3 Products) |

Target Details

| Bac | kar | 'nΙ | ın | Ч. |
|-----|-----|--------|----------|----|
| Duo | | \sim | <i>.</i> | ч. |

Synonyms: cAMP responsive element binding protein 3 like 3, cAMP responsive element-binding protein 3 like protein 3, CREB H, CREB/ATF family transcription factor, CREBH, Cyclic AMP responsive element binding protein 3 like protein 3, cAMP-responsive element-binding protein 3-like protein 3, HYST1481, MGC126553, MGC126557, Transcription factor CREB H, CR3L3_HUMAN.

Background: CREB3L3 is a 461 amino acid single-pass type II membrane protein that localizes to the endoplasmic reticulum (ER) and, in response to ER stress, is cleaved and translocated to the nucleus. Expressed exclusively in liver, CREB3L3 functions as a transcription factor that, during ER stress, is thought to activate genes that are involved in both the unfolded protein response and the acute phase response (APR). Additionally, CREB3L3 is underexpressed in hepatocellular carcinoma, suggesting a possible role as a tumor suppressor. CREB3L3 functions as a dimer and contains one leucine zipper domain, a KDEL-like sequence and a bZIP domain, through which it conveys its DNA binding ability. Three isoforms of CREB3L3 exist due to alternative splicing events.

Pathways:

Thyroid Hormone Synthesis

Application Details

| Δnn | lication | Notae. |
|-----|----------|--------|
| | | |

FCM 1:20-100

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |