

# Datasheet for ABIN2811928

# anti-CRH antibody (AA 185-196) (AbBy Fluor® 594)



Go to Product page

Overview	
Quantity:	100 μL
Target:	CRH
Binding Specificity:	AA 185-196
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRH antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CRF
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Pig,Chicken,Rabbit,Guinea Pig
Purification:	Purified by Protein A.
Target Details	
Target:	CRH
Alternative Name:	CRF (CRH Products)

# Target Details

Bac	kar	OL.	ın	d:
Duo		$\sim$		u.

Synonyms: Corticoliberin, Corticoliberin precursor, Corticotropin releasing factor, Corticotropin releasing hormone, Corticotropin releasing hormone deficiency included, crf, crh, crh deficiency included, corticoliberin preproprotein, CRF\_HUMAN, Corticotropin-releasing factor, CRF, CRH, Corticotropin-releasing hormone.

Background: Corticotropin-releasing hormone is secreted by the paraventricular nucleus (PVN) of the hypothalamus in response to stress. Marked reduction in this protein has been observed in association with Alzheimer disease and autosomal recessive hypothalamic corticotropin deficiency has multiple and potentially fatal metabolic consequences including hypoglycemia and hepatitis. In addition to production in the hypothalamus, this protein is also synthesized in peripheral tissues, such as T lymphocytes and is highly expressed in the placenta. In the placenta it is a marker that determines the length of gestation and the timing of parturition and delivery. A rapid increase in circulating levels of the hormone occurs at the onset of parturition, suggesting that, in addition to its metabolic functions, this protein may act as a trigger for parturition. [provided by RefSeq, Apr 2010].

Gene	11 ).
Oene	ID.

1392

### UniProt:

P06850

#### Pathways:

Positive Regulation of Peptide Hormone Secretion, Hormone Activity, Negative Regulation of Hormone Secretion, cAMP Metabolic Process, Myometrial Relaxation and Contraction, Feeding Behaviour

## **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

# Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months