antibodies .- online.com





anti-CRTC3 antibody (AA 151-260) (Alexa Fluor 594)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	CRTC3
Binding Specificity:	AA 151-260
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRTC3 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRTC3/TORC3	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	CRTC3
Alternative Name:	CRTC3/TORC3 (CRTC3 Products)
Background:	Synonyms: CREB regulated transcription coactivator 3, CREB-regulated transcription

coactivator 3, CRTC 3, CRTC3, CRTC3_HUMAN, FLJ21868, TORC 3, TORC-3, TORC3,
Transducer of CREB protein 3, Transducer of regulated cAMP response element binding protein
CREB 3, Transducer of regulated cAMP response element binding protein 3, Transducer of
regulated cAMP response element-binding protein 3, Transducer of regulated CREB protein 3.
Background: TORC3 is a 619 amino acid protein that localizes to both the cytoplasm and the
nucleus and belongs to the TORC family. Expressed in lung tissue and B and T lymphocytes, as
well as in colon, brain, ovary, kidney, prostate, colon and heart, TORC3 functions as a
transcriptional coactivator for CREB-1, thereby regulating the expression of CREB-activated
genes, and is also thought to activate the SIK/TORC signaling pathway. TORC3 exists as
multiple alternatively spliced isoforms and, in addition to its role in transcriptional activation, is
thought to induce mitochondrial biogenesis, specifically in muscle cells.

Gene ID: 64784

Pathways: SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2

Infection

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

ш	Jand	lina
г	land	1111()
•	10110	9

Expiry Date:

12 months