# antibodies .- online.com





# anti-CSAD antibody (AA 401-493) (Alexa Fluor 594)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | CSAD   |
| Binding Specificity: | AA 401-493   |
| Reactivity:          | Mouse, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This CSAD antibody is conjugated to Alexa Fluor 594  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human CSAD |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse, Rat   |
| Predicted Reactivity: | Human,Dog,Horse  |
| Purification:         | Purified by Protein A.                                   |

## **Target Details**

| Target:           | CSAD                 |
|-------------------|----------------------|
| Alternative Name: | CSAD (CSAD Products) |

## **Target Details**

| Background:         | Synonyms: CSAD, CSAD_HUMAN, CSD, Cysteine sulfinic acid decarboxylase, Cysteine-sulfinate   |  |
|---------------------|---|--|
|                     | decarboxylase, Sulfinoalanine decarboxylase.  |  |
|                     | Background: CSAD is a 493 amino acid protein that exists as a homodimer and belongs to the  |  |
|                     | group II decarboxylase family. CSAD catalyzes the conversion of 3-sulfino-L-alanine to      |  |
|                     | hypotaurine and carbon dioxide, binds pyridoxal phosphate as a cofactor and undergoes       |  |
|                     | alternative splicing to produce three isoforms. The gene encoding CSAD maps to human        |  |
|                     | chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5 $\%$ of the   |  |
|                     | human genome. Chromosome 12 is associated with a variety of diseases and afflictions,       |  |
|                     | including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and        |  |
|                     | trisomy 12p, which causes facial developmental defects and seizure disorders.               |  |
| Gene ID:            | 51380   |  |
| Application Details |   |  |
| Application Notes:  | IF(IHC-P) 1:50-200  |  |
|                     | IF(IHC-F) 1:50-200  |  |
|                     | IF(ICC) 1:50-200  |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 μg/μL   |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |  |
|                     | 50 % Glycerol.  |  |
| Preservative:       | ProClin   |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be         |  |
|                     | handled by trained staff only.  |  |
| Storage:            | -20 °C  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.           |  |
| Expiry Date:        | 12 months   |  |