antibodies .- online.com

Datasheet for ABIN2812053 anti-CUEDC1 antibody (AA 251-350) (Alexa Fluor 594)



Overview

Quantity:	100 µL
Target:	CUEDC1
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CUEDC1 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CUEDC1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	CUEDC1
Alternative Name:	CUEDC1 (CUEDC1 Products)
Background:	Synonyms: CUE domain containing protein 1, CUED1_HUMAN.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2812053 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

(CUE) domain functions as a ubiquitin (UB) binding domain that is approximately 40 ar acids in length. Present in eukaryotic proteins that are involved in ubiquitination and pr trafficking pathways, CUE domains can bind monoubiquitin and may be required for ubiquitination of the proteins in which they are found. CUEDC1 (CUE domain-containin 1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible ro protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 m associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor r Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Buffer: Liquid Concentration: 1 µg/µL </th <th></th> <th></th>		
acids in length. Present in eukayotic proteins that are involved in ubiquitination and pr trafficking pathways, CUE domains can bind monoubiquitin and may be required for ubiquitination of the proteins in which they are found. CUEDC1 (CUE domain-containin 1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible ro protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 in associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor in Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Handling Image: Concentration: Handling Image: Concentration: Handling Image: Concentration: Precentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procheed 50 % Glycerol.		Background: The coupling of ubiquitin conjugation to endoplasmic reticulum (ER) degradation
trafficking pathways, CUE domains can bind monoubiquitin and may be required for ubiquitination of the proteins in which they are found. CUEDC1 (CUE domain-containin 1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible ro protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 massociated with early stage cervical cancer, implicating CUEDC1 as a potential tumor r Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 IF(UCC) 1:50-200 Restrictions: For Research Use only Handling Eormat: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclisto % Glycerol. Preservative: ProClin Preservative: <		(CUE) domain functions as a ubiquitin (UB) binding domain that is approximately 40 amino
ubiquitination of the proteins in which they are found. CUEDC1 (CUE domain-containin 1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible ro protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 in associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor in Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200		acids in length. Present in eukaryotic proteins that are involved in ubiquitination and protein
1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible ro protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 massociated with early stage cervical cancer, implicating CUEDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 as a potential turnor massociated with early stage cervical cancer, implicating cueDC1 exist due to alternative splicing events. Application Details IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 Handling Early content to a content on the only If and the only Handling Iuguid Iuguid Iuguid Concent		trafficking pathways, CUE domains can bind monoubiquitin and may be required for
protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 as a potential tumor r associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor r Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		ubiquitination of the proteins in which they are found. CUEDC1 (CUE domain-containing protein
associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor r Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		1) is a 386 amino acid protein that contains one CUE domain, suggesting a possible role in
Two isoforms of CUEDC1 exist due to alternative splicing events. Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procision % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		protein trafficking and degradation pathways. Defects in the gene encoding CUEDC1 may be
Gene ID: 404093 Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclises 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		associated with early stage cervical cancer, implicating CUEDC1 as a potential tumor marker.
Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Implication Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procelist 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		Two isoforms of CUEDC1 exist due to alternative splicing events.
Application Notes: IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Iuid Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclis 50 % Glycerol. Preservative: Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Gene ID:	404093
IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procells 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Details	
IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which shandled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Notes:	IF(IHC-P) 1:50-200
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procling 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which is handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(IHC-F) 1:50-200
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclit 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(ICC) 1:50-200
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only
Concentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Procli 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Handling	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclis 50 % Glycerol. Storage: Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Format:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Concentration:	1 μg/μL
Precaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which s handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Preservative:	ProClin
Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		handled by trained staff only.
	Storage:	-20 °C
Expiry Date: 12 months	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
	Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN2812053 | 03/07/2024 | Copyright antibodies-online. All rights reserved.