# antibodies .- online.com





# anti-CWC22 antibody (AA 101-200) (AbBy Fluor® 594)



$C_0 + 0$	Product	0000
(¬() (()	Promici	Dage

$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	CWC22
Binding Specificity:	AA 101-200
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CWC22 antibody is conjugated to AbBy Fluor® 594
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CWC22
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	CWC22
Alternative Name:	CWC22 (CWC22 Products)

## **Target Details**

Background:	Synonyms: CWC22, CWC22 spliceosome associated protein homolog S. cerevisiae, CWC22
	spliceosome associated protein homolog, CWC22_HUMAN, E4GL, fSAPb, Functional
	spliceosome associated protein b, KIAA1604, NCM, Nucampholin homolog, Pre-mRNA-splicing
	factor CWC22 homolog.
	Background: Spliceosomes are multi-protein complexes that are composed of snRNPs (small
	nuclear ribonucleoproteins) and a variety of associated protein factors, all of which work in
	concert to regulate the splicing of pre-mRNA, a critical step in the post-transcriptional
	regulation of gene expression. CWC22 (CWC22 spliceosome-associated protein), also known
	as NCM, fSAPb or EIF4GL, is a 908 amino acid nuclear protein and component of the
	spliceosome C complex. CWC22 is associated with the spliceosome prior to catalytic steps and
	remains associated throughout the reaction. Belonging to the CWC22 family, CW22 contains
	one MI domain and a MIF4G domain. The gene encoding CWC22 maps to human chromosome
	2q31.3. As the second largest human chromosome, chromosome 2 makes up approximately
	8 % of the human genome and contains 237 million bases encoding over 1,400 genes.
Gene ID:	57703
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months