# antibodies -online.com





# anti-CYP2R1 antibody (AA 251-350) (AbBy Fluor® 594)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	CYP2R1
Binding Specificity:	AA 251-350
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP2R1 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CYP2R1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	CYP2R1
Alternative Name:	CYP2R1 (CYP2R1 Products)

### **Target Details**

Background:	Synonyms: Cytochrome P450 2R1, Cytochrome P450 family 2 R1, Cytochrome P450 family 2	
	subfamily R polypeptide 1, EC 1.14.13.15, MGC4663, Vitamin D 25 hydroxylase.	
	Background: CYP2R1 is a member of the cytochrome P450 superfamily of enzymes. The	
	cytochrome P450 proteins are mono-oxygenases which catalyze many reactions involved in	
	drug metabolism and synthesis of cholesterol, steroids and other lipids. This enzyme is a	
	microsomal vitamin D hydroxylase that converts vitamin D into the active ligand for the vitamin	
	D receptor.Defects in CYP2R1 are a cause of 25-hydroxyvitamin D(3) deficiency, also known as	
	pseudovitamin D(3) deficiency rickets due to 25-hydroxylase deficiency. First described in	
	patients who had rickets at a young age despite a history of adequate vitamin D intake. The	
	patients sera had low calcium concentrations, low phosphate concentrations, elevated alkaline	
	phosphatase activity and low levels of 25-hydroxyvitamin D.	
Gene ID:	120227	
Pathways:	Metabolism of Steroid Hormones and Vitamin D	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	