

[Go to Product page](#)

## Datasheet for ABIN2812177 **anti-CST4 antibody (Alexa Fluor 594)**

### Overview

Quantity:	100 µL
Target:	CST4
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CST4 antibody is conjugated to Alexa Fluor 594
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cystatin S
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

### Target Details

Target:	CST4
Alternative Name:	Cystatin S ( <a href="#">CST4 Products</a> )
Background:	<p>Synonyms: Cst4, Cystatin 4, Cystatin S precursor, Cystatin SA III, Cystatin-4, Cystatin-S, Cystatin-SA-III, CYTS_HUMAN, MGC71923, Salivary acidic protein 1.</p> <p>Background: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost</p>

Target Details

or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins(stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a type 2 salivary cysteine peptidase inhibitor. The protein is an S-type cystatin, based on its high level of expression in saliva, tears and seminal plasma. The specific role in these fluids is unclear but antibacterial and antiviral activity is present, consistent with a protective function. [provided by RefSeq, Jul2008].

Gene ID: 1472

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months